

# On locally invertible harmonic mappings of plane domains

Avkhadiev F.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017, Pleiades Publishing, Ltd. We prove several global univalence theorems for locally invertible harmonic mappings with certain prescribed boundary behavior in simply and multiply connected domains. In particular, we consider mappings with singular boundary points when the argument principle is not applicable. In addition, we examine harmonic mappings connected with the famous von Mises coordinates. It is shown the univalence of every locally univalent von Mises harmonic mapping defined in a domain convex in the direction of ordinate axis.

<http://dx.doi.org/10.1134/S1995080217030076>

---

## Keywords

argument principle, Hadamard's theorem, Harmonic mapping, local homeomorphism, Mises coordinates

## References

- [1] A. G. Sergeev, Harmonic Maps, Vol. 10 of Lect. Notes SEC (Inst. Mat. RAN Steklova, Moscow, 2008) [in Russian].
- [2] V. N. Monakhov, Boundary-Value Problems with Free Boundaries for Elliptic Systems of Equations (Am. Math. Soc., Providence, RI, 1983).
- [3] M. H. Hamdan and R. M. Barron, "Applications of von Mises coordinates in porous media flow," J. Comput. Appl. Math. 39, 353–361 (1996).
- [4] H. Lewy, "On the non-vanishing of the Jacobian in certain one-to-one mappings," Bull. Am. Math. Soc. 42, 689–692 (1936).
- [5] P. Duren, Harmonic Mappings in the Plane (Cambridge Univ. Press, Cambridge, 2004).
- [6] M. Morse, Topological Methods in the Theory of Functions of a Complex Variable Vol. 15 of Ann. Math. Studies (Princeton Univ. Press, New York, 1947).
- [7] F. G. Avkhadiev, "On injectivities in domains of open isolated mappings with given boundary properties," Sov. Math. Dokl. 35, 117–120 (1987).
- [8] J. Hadamard, "Sur les transformations ponctuelles," Bull. Soc. Math. France 34, 71–84 (1906).
- [9] S. Banach and S. Masur, "Über mehrdeutige stetige Abbildungen," Studia Math. 5, 174–178 (1934).
- [10] F. E. Browder, "Covering spaces, fiber spaces and local homeomorphisms," Duke Math. J. 21, 329–336 (1954).
- [11] T. Parthasarathy, On Global Univalence Theorems (Springer, Berlin, Heidelberg, New York, 1983).
- [12] F. G. Avkhadiev and K.-J. Wirths, "Conformal radius as a function and its gradient images," Israel. J. Math. 145, 349–374 (2005).
- [13] F. G. Avkhadiev and K.-J. Wirths, Schwarz–Pick Type Inequalities (Birkhäuser, Basel, Boston, Berlin, 2009).
- [14] P. S. Alexandrov, Combinatorial Topology, Dover Books in Math. (Dover, New York, 1998).

- [15] F. G. Avkhadiev, "Sufficient conditions for the univalence of quasiconformal mappings," Math. Notes 18, 1063–1067 (1975).
- [16] T. Umezawa, "On the theory of univalent functions," Tohoku Math. J. 5, 218–228 (1955).
- [17] M. Chuaqui and R. Hernández, "Univalent harmonic mappings and linearly connected domains," J. Math. Anal. Appl. 332, 1189–1194 (2007).